


## PAT-122 4/00





Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99203197.1

TECHNOLOGY CENTER 1700

RECEIVED

RECEIVED

JAN 5 2001

TC 1700

Der Präsident des Europäischen Patentamts;  
Im Auftrag


For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE,  
LA HAYE, LE

27/09/00





Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.: 99203197.1  
Demande n°:

Anmeldetag:  
Date of filing: 30/09/99  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
**QUEST INTERNATIONAL B.V.**  
**1411 GP Naarden**  
**NETHERLANDS**

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

**Savoury flavour comprising 2-methyl-furan -3-thiol and/or a derivative and methylenedithiol and/or a derivative**

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

**A23L1/226, C07C319/02**

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:



Savoury flavour comprising 2-methylfuran-3-thiol and/or a derivative and methylenedithiol and/or a derivative.

5

The present invention relates to flavoured foodstuffs and flavouring compositions. More in particular the present invention relates to foodstuffs having a savoury flavour and to flavouring compositions suitable for imparting such  
10 a flavour. Under a savoury flavour is here to be understood a flavour associated with meat, sausage, poultry, cheese etc. The term foodstuff as used herein includes both solid and liquid ingestible materials which may or may not have nutritional value.

15

Flavoured foodstuffs having a savoury flavour and flavouring compositions for imparting to and/or enhancing in a foodstuffs a savoury flavour have been known for a long time, but so far these compositions were not been  
20 quite satisfactory. Therefore the present invention aims to improve the organoleptic properties.

Many compounds have been used to impart a savoury flavour to foodstuffs and therefore most of the compositions used  
25 for imparting a savoury flavour contain a multitude of compounds. One group of particular useful compounds for savoury flavours are certain (hydrogenated) furan thiols and derivatives thereof like thio-acetates are e.g. disclosed and/or claimed in GB-A-1 283 912 (Unilever). GB-  
30 A-1 256 462 (International Flavours and Fragrances) also discloses certain furan thiols such as 2-methylfuran-3-thiol and its derivatives such as its disulphide and other sulphides as meat flavours. Especially 2-methylfuran-3-thiol and its precursors are powerful flavouring compounds  
35 capable of imparting a pleasant roast meat flavour to foodstuffs [cf e.g. "Volatile compounds in foods en beverages" by H. Maarse (ED) Marcel Dekker, New York (1991)]. Its powerful roast or fried beef note is, however, not always


BO 43104

2

appreciated. Therefore attempts have been made to obtain products with a powerful, somewhat milder, more rounded off meat flavour more reminiscent of the flavour of boiled beef  
5 and beef broth.

The present invention solves the problem of obtaining a powerful mild beef flavour reminiscent of beef broth, and boiled beef which is especially needed for soups, sausages,  
10 pastry etc. Also varying the relative amounts of the key components the present invention enables the flavourist to prepare flavouring compositions varying from a mild roast beef flavour to a mild beef broth flavour.

15 In a first embodiment the present invention therefore provides a flavoured foodstuff comprising an effective amount of at least one compound with a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an -S-CH<sub>3</sub> group, an -CO-CH<sub>3</sub> group or a 2-methyl-3-furyl-thio moiety  
20 and an effective amount of at least one compound having the structure



25 in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a  
30 lower acyl group or a 2-methyl-3-furyl-thio group or a -S-CH<sub>2</sub>-U group as defined above.

Suitable compounds in the practice of the present invention having a (hydrogenated) 2-methyl-3-furyl-thio moiety and a  
35 hydrogen atom, an -S-CH<sub>3</sub> group, an -CO-CH<sub>3</sub> group or a 2-methyl-3-furyl-thio moiety are e.g. 2-methylfuran-3-thiol,




BO 43104

3

2-methyl-4,5-dihydrofuran-3-thiol, cis/trans 2-methyltetrahydrofuran-3-thiol, 2-methyl-3-thiomethoxyfuran, methyl-2-methyl-3-furyldisulphide, 2-methylfuran-3-thioacetate, 2-methylfuran-3-thiopropionate and the disulphide of 2-methylfuran-3-thiol. The use of non-hydrogenated furanderivatives is preferred. Several of these compounds are available from Oxford Chemicals, Hartlepool, U.K. It is observed that 2-methylfuran-3-thiol like many other thiols is liable to oxidise partially under the influence of oxygen in the atmosphere to its disulphide and consequently quite often commercially available 2-methylfuran-3-thiol preparations contain certain amounts of its disulphide. Thioesters like 2-methyl-3-furan-3-thioacetate and e.g. 2-methyl-3-furan-3-thiopropionate may hydrolyse partially to form 2-methyl-3-furan-3-thiol under certain food processing conditions. This may apply e.g. under the conventional sterilising conditions after canning.

Suitable compounds for the practice of the present invention having a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a 2-methyl-3-furyl-thio moiety and an effective amount of at least one compound having the structure



in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a lower acyl group or a 2-methyl-3-furyl-thio group or a  $-S-CH_2-U$  group as defined above are methanedithiol, methanedithiol diacetate, methylthiomethanol, methylthio-

BO 43104

4

methanethiol, methylthiomethanethiol acetate, methylthio-  
methanethiol propionate, methylthiomethanethiol 2-  
methylpropionate, methylthiomethanethiol 2-methylbutanoate,  
5 methylthiomethanethiol 2-methylbutanoate, 3-methyl-  
thiomethanethiol pentanoate, methylthiomethanethiol 4-  
methylpentanoate and methylthiomethanethiolhexanoate. Some  
of these compounds are to be regarded as precursors of the  
corresponding free thiols, whereas others have organoleptic  
10 properties resembling methanedithiol or methylthiomethanol  
etc. with slightly different flavour notes. Here also the  
possibility arises that free thiols are partially converted  
to corresponding disulphides under oxidising conditions and  
also thioesters may hydrolyse to form free thiols under  
15 certain food processing conditions.

The present invention not only covers foodstuffs flavoured  
by incorporating the above identified compounds per se, but  
also foodstuffs incorporating precursors of the above  
20 identified compounds.

2-Methyl-3-furanthiol and some of its derivatives discussed  
above can be obtained or synthesized as such, but it may  
also be convenient to prepare a meat/beef flavour  
25 composition containing one or more of these compounds (a  
reaction flavour composition) which comprises at least one  
of these compounds in a suitable quantity and incorporate  
in such a composition a suitable amount of a compound  
having the structure  $U-CH_2-S-T$ . Suitable reaction  
30 flavour compositions can be prepared by reacting a hexose  
or pentose with a source of hydrogen sulphide such as  
cysteine in water as a solvent. Instead of a hexose or  
pentose a suitable degradation product of a sugar can be  
used such as e.g. 4-hydroxy-5-methyl-2,3-hydrofuran-3-one  
35 and ascorbic acid may also be used. It is also quite  
possible to prepare suitable reaction flavours containing  
2-methyl-3-furanthiol by thermal degradation of thiamine.

BO 43104

5

Several possibilities and reaction mechanisms of reaction flavours are discussed by Hoffmann & Schieberle in J. Agric. Food Chemistry, 46, 235-241 (1998) which is hereby  
5 incorporated by reference.

The flavouring compositions according to the present invention are especially incorporated in foodstuffs such as sausage, meat, reformed comminuted meat, meat replacers  
10 such as texturized vegetable protein, and pastry products in an effective amount i.e. in an amount sufficient to impart or reinforce the desired flavour. As a general guide flavouring amounts of 0.01 to 1000, preferably 0.5 to 100  
15 ppb (parts per billion =  $1 : 10^9$ ) are incorporated in foodstuffs which proportions are expressed as on a weight basis calculated on the foodstuff. The flavouring amounts actually incorporated depend on the individual palate and  
20 on the nature of the foodstuff. In flavouring compositions the amounts present may easily be a factor 10 to 10.000 higher than in the actual flavoured foodstuff.

Preferably the present invention provides a flavoured foodstuff as defined above, in which lower thioacyl- and lower acyl group means that these groups comprise from 2 to  
25 6, preferably 2 or 3 carbon atoms.

Preferably the present invention provides a flavoured foodstuff as defined above, in which U represents a lower thioacyl group and T represents a lower acyl group.  
30


More preferably the present invention provides a flavoured foodstuff as defined above, in which lower thioacyl group means thioacetoxyl and lower acyl group independently means acetyl.  
35

In another embodiment the present invention provides a process for imparting a savoury flavour to a foodstuff

BO 43104

6

comprising incorporating in said foodstuff an effective amount of at least one compound with a 2-methyl-3-furyl-thio moiety and a hydrogen atom, an -S-CH<sub>3</sub> group, an -CO-CH<sub>3</sub> group or a (hydrogenated) 2-methyl-3-furyl moiety and an effective amount of at least one compound having the structure



10

in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a lower acyl group or a 2-methyl-3-furyl-thio moiety or a -S-CH<sub>2</sub>-U group as defined above.


Preferably the above process according to the present invention employs compounds as defined above, in which lower thioacyl-, lower alkyl- and lower acyl group means that these groups comprise from 2 to 6, preferably 2 or 3 carbon atoms.

More preferably the above process according to the present invention employs at least one compound as defined above, in which U represents a lower thioacyl group or a lower acyloxy group and T represents a lower acyl group.

In an other embodiment the present invention provides a flavouring composition for foodstuffs comprising at least one compound comprising a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an -S-CH<sub>3</sub> group, an -CO-CH<sub>3</sub> group or a (hydrogenated) 2-methyl-3-furyl-thio moiety and at least one compound having the structure

BO 43104

7



in which C, H and S have the conventional meanings of  
5 carbon, hydrogen and sulphur atoms respectively, U  
represents a thiol group, a lower thioacyl group, a  
hydroxyl group or a 2-methyl-3-furyldithiogroup and T  
represents a hydrogen atom, a lower alkyl group or a lower  
acyl group. Such a flavouring composition can be used for  
10 enhancing an existing weak savoury flavour e.g. to  
compensate for flavour notes lost by processing foodstuffs,  
but they can equally be used for flavouring a bland or  
tasteless foodstuff. Moreover it is also possible to change  
the flavour characteristics of a foodstuff completely.  
15 Flavouring compositions are frequently available in the  
form of active material in a suspension or a solution or  
upon an organoleptically inactive material.

Preferably the flavouring composition for foodstuffs  
20 according to the present invention comprises at least one  
or more compounds as defined above, in which lower  
thioacyl-, lower alkyl- and lower acyl group means that  
these groups comprise from 2 to 6, preferably 2 or 3 carbon  
atoms.

25

The flavouring compositions for foodstuffs may be in liquid  
or semi-liquid form such as solutions, emulsions or pastes,  
or in dry form such as a powder. Drying can be accomplished  
for example by spray-drying or by freeze-drying, optionally  
30 on a carrier such as maltodextrin.


The present invention is not restricted to at least one  
compound with a (hydrogenated) 2-methyl-3-furanthiol  
(derivative) together with at least one compound with a  
35 methylenedithiol (derivative) as defined above. As is  
common in flavour creation other compounds known to  
contribute to a savoury flavour can be incorporated as

BO 43104

8

well. Such known compounds are e.g. amino acids, nucleotides, monosodiumglutamate, lower alcohols, lower carboxylic acids, pyrrolidone carboxylic acid, lower peptides, sweeteners, lactones, lower disulphides, lower thiols, guanidines etc., salts like NaCl, amines, lower aldehydes, lower ketones, tricholomic acid, biotenic acid, aromatic and/or heterocyclic compounds like acetyl thiazole, 2-hydroxyethyl-4-methylthiazol, 4-hydroxy-2,5-dimethyl-2,3-hydrofuran-3-one, colouring materials, thickening agents. The proportions of these optionally added substances used are dependent on the kind of flavour desired and also on the nature of the foodstuffs in which they are incorporated and also on any herbs or spices added.

According to another embodiment the present invention provides the use of both an effective amount of at least one compound with a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a 2-methyl-3-furyl-thio group and an effective amount of at least one compound having the structure



25

in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a lower acyl group or a 2-methyl-3-furyl-thio group or a  $-S-CH_2-U$  group as defined above.

According to another embodiment of the present invention provides a process for preparing a pure compound with at least one free thiol group as defined in claim 1 by

BO 43104

9

hydrolysing the corresponding thioacyl compound in the presence of an enzyme or a cation exchange resin.

- 5 More preferably the process defined above comprises the use of a lipase. More in particular lipases like lipase 435 ex NOVO, Denmark can be used with good results.

The invention will be illustrated by the following examples  
10 in which all parts and percentages etc. are on a weight basis unless otherwise indicated.

#### Example 1

##### Methanedithiol diacetate

- 15 Diiodomethane (26.8 g, 0.1 mole) was dosed to a mixture of 250 ml acetone, potassium thioacetate (30 g) and molecular sieves 3A (5 g) over 20 minutes at 20°C. The reaction temperature rose to 38°C (without external cooling). The mixture was stirred during 2 hours at room temperature and  
20 quenched in 500 ml water. The resulting mixture was extracted three times with 100 ml methyl tertiary butylether (MTBE) each. The MTBE extracts were joined, washed twice with 100 ml water and dried over anhydrous magnesium sulphate. The solvent was evaporated using a  
25 rotavapour (water bath 30°C at 8 Kpa (80 mbar). Obtained were 18 g concentrate. The concentrate was distilled using a 10 cm Vigreux column. Obtained were 13.5 g distillate with a boiling point of 86°C at 0.3 Kpa (3 mbar). According to gas chromatography the distillate was more than 99% pure  
30 and its NMR data matched. Yield 85% of the theoretical yield.

#### Example 2

##### Methanedithiol

- 35 Methanedithiol diacetate (10 g) was dissolved in methanol (80 g) containing 2 g of gaseous hydrochloric acid and the mixture was stirred overnight at room temperature. The solution was quenched in 200 ml water and the obtained

BO 43104

10

solution was three times extracted with each 50 ml dichloromethane. The dichloromethane extracts were joined, washed with 50 ml water and dried over anhydrous magnesium sulphate. The dichloromethane was evaporated using a 10 cm Vigreux column until a bottom temperature of 20°C at 4.3 Kpa (43 mbar) was reached. Obtained were 8 g concentrate. The concentrate was distilled with the same apparatus. Obtained were 3.5 g distillate with a boiling point of 35°C at 4.3 Kpa. The NMR data matched and the yield was 70% of the theoretical yield. The distillate was a suitable ingredient for a savoury flavour composition according to the invention.

### 15 Example 3

#### Methanedithiol

Methanedithiol diacetate (4.9 mole, 0.80 g) obtained as described above was taken up in pentane (5 g) as solvent and transesterified with ethanol (10.9 mmol) and lipase SP 20 435 ex NOVO as catalyst. The ethanol was slowly added over 12 hours. The total reaction time was 12 hours at 20°C. Gas chromatographic analysis indicated that the conversion was quantitative. Subsequently the enzyme immobilisate was filtered off and the pentane solution concentrated. The 25 reaction mixture was analyzed with NMR and matched. The results are tabulated below:

Substance	Molar ratio
Ethyl acetate	9
30 Methanedithiol	8

The concentrate was diluted with ethanol and found to contribute to form an excellent cooked meat flavour especially in combination with 2-methyl-furan-3-thiol.

35



BO 43104

11

Example 4**Methylthiomethanethiol acetate**

Chloromethylmethylsulphide (4.8 g, 50 mmole) in methyl  
5 tert.-butylether (50 mL) as solvent was treated at 10-12°C  
with thioacetic acid (4.0 g, 50 mmole), dissolved in methyl  
tert. butylether (2 mL). Subsequently triethylamine (5.1 g,  
50 mmole) in methyl tert. butyl ether (2 mL) were added to  
the reaction mixture at 14-16°C. The reaction mixture was  
10 stirred for 3 hours at 20°C. Water (200 mL) was added and  
the mixture was stirred for 15 minutes. The organic layer  
was separated, and the aqueous layer was extracted twice  
with methyl tert. butyl ether. The combined extracts were  
dried over anhydrous magnesium sulphate, filtered and  
15 concentrated. The residue was distilled and fractionated  
under reduced pressure. The results are tabulated below:

Fraction	Bottom temperature	Top temperature	Pressure	Amount
1	97°C	96°C	5.4 Kpa	0.2 g
2	105°C	104°C	5.4 Kpa	0.9 g
3	112°C	105°C	5.4 Kpa	3.2 g

Fractions 2 and 3 were satisfactory for flavouring  
purposes.

Example 5**Methylthiomethanethiol**

A mixture of methylthiomethanethiol acetate (2.04 g, 15  
mmole) and a 15% aqueous sodium hydroxide solution (14.8 g)  
30 was stirred for 2 hours at 75°C. The mixture was cooled to  
0-5°C and a 10% aqueous Hcl solution (21.7 g) was slowly  
added at 0-5°C. The mixture was extracted with ether three  
times, and the combined ether extracts were dried over  
anhydrous magnesium sulphate. After filtration and

BO 43104

12

concentration of the filtrate the residue was distilled at 75°C at 8.0 Kpa (60 Torr) to give 0.19 g of the pure title compound and was a suitable ingredient for flavouring 5 compositions.

Example 6

Methylthiomethanethiol.

Methylthiomethanethiol acetate was transesterified with 10 ethanol by the method described in Example 3 above to yield a solution of methylthiomethanethiol in ethanol. As indicated by gas chromatographic analysis the conversion into methylthiomethanethiol was complete after 12 hours. The solution was satisfactory for flavouring purposes.

15

Example 7

Methylthiomethanethiol

Sodium sulphide hydrate (0.25 g) was covered with dichloromethane. A buffer solution of Ph 5.3 (NaOAc/HOAc) 20 (82 g sodium acetate and 25.3 g acetic acid dissolved in water, total volume 711 Ml) was carefully poured on the dichloromethane, followed by formaldehyde 37% in water (20.2 g, 0.25 mmol). The bottle was purged with nitrogen and closed with a screw cap with a silicon rubber septum. 25 The mixture was stirred magnetically for three days. The aqueous layer was extracted with dichloromethane. The crude product, in dichloromethane as solvent had a composition as tabulated below:

30	<table><tr><th>Substance</th><th>Molar ratio</th></tr><tr><td>Methylthiomethanethiol</td><td>3-4</td></tr><tr><td>Methanethiol</td><td>2</td></tr><tr><td>Methylthiomethanol</td><td>1</td></tr></table>	Substance	Molar ratio	Methylthiomethanethiol	3-4	Methanethiol	2	Methylthiomethanol	1
Substance	Molar ratio								
Methylthiomethanethiol	3-4								
Methanethiol	2								
Methylthiomethanol	1								

35

BO 43104

13

The solvent was removed under reduced pressure and the product was distilled twice under reduced pressure to yield a product fraction which distilled at 45-47°C at 2 Kpa (20 5 mbar). The product fraction, which was a suitable meat flavour component, had a composition as tabulated below.

10	Substance	Molar ratio
	Methylthiomethanethiol	16.5
	Methanethiol	2.4
	Methylthiomethanol	1

The three main components are responsible for 75% of the 15 integral in the NMR.

#### Example 8

##### Various disulphides

2-methylfuran-3-thiol (10 mg) and methylthiomethanethiol 20 (10 mg) were taken up in 1 mL ethanol in the presence of basic aluminium trioxide. The mixture was stirred for 20 hours at 20°C under a slight overpressure of oxygen. Subsequently the aluminium trioxide was removed from the solution by centrifuging. The centrifuged solution was a 25 valuable ingredient for savoury flavours. The solution was analyzed by GC/MS and the results are tabulated below.

BO 43104

14

5	Compound	%	GC/MS data
	2-Methylfuran-3-thiol	trace	
10	Methylthiomethanethiol	trace	
	Bis(2-methylfuran-3-)-disulphide	29%	226(79), 162(9), 155(7), 113(110), 85(15), 43(9)
	Bis(methylthiomethane)-disulphide	29%	186(29), 153(6), 93(8), 61(100), 45(15)
	(2-Methylfuran-3)-methylthiomethanedisulphide	31%	206(40), 113(39), 95(48), 93(11), 61(100), 45(14)

15

Example 9

A reaction flavour block with flavour type meaty/beefy was prepared by the thermal reaction of xylose with cysteine hydrochloride in water as the solvent, followed by spray-drying the reaction mixture on a maltodextrin 10 support. The spray-dried flavouring composition contained 2-methyl-3-furanthiol at a concentration of 20 ppm. An amount of 0.5 g of the spray-dried beef flavour was dissolved in 1 L water with a temperature of 50°C containing sodium chloride (5 g). The resulting solution (test solution A) contained 2-methyl-3-furanthiol at a concentration of 10 ppb (parts per billion). To a 0.5 L aliquot of mixture A were added 25 microlitres of a solution of methylthiomethanethiol in water (200 ppm). The obtained solution (test solution B) contained 2-methyl-3-furanthiol at a concentration of 10 ppb and methylthiomethanethiol at a concentration of 10

BO 43104

15

ppb. The flavoured test solutions were evaluated by a panel of four experienced tasters. Three out of the four tasters found test solution B to possess a distinctly cooked note, 5 which was absent in test solution A.

#### Example 10

Four savoury flavour compositions were prepared by mixing 10 the following ingredients in the amounts (parts per thousand) indicated in the table below:

The four mixtures A, B, C and D so obtained were added separately to a test solution (50°C, containing 5 g/L 15 sodium chloride) at a level of 0.04 g per litre. The flavoured test solutions were evaluated by a panel of four experienced tasters. Three of the four tasters described both mixtures C and D as having an enhanced cooked note in comparison with mixtures A and B. Three experienced tasters 20 were of the opinion that both mixtures C and D displayed a less brown and roasted note than mixture B.

BO 43104

16

	Compound	Mixt. A	Mixt. B	Mixt. C	Mixt. D
	Hexanal	11.4	11.4	11.4	11.4
	Nonanal	2.0	2.0	2.0	2.0
5	t-2-Heptenal	1.2	1.2	1.2	1.2
	t-2-Octenal	1.2	1.2	1.2	1.2
	t-2-Nonenal	1.2	1.2	1.2	1.2
	t-2-Decenal	3.5	3.5	3.6	3.5
	t,t-2,4-Heptadienal	1.0	1.0	1.0	1.0
10	t,t-2,4-Nonadienal	0.8	0.8	0.8	0.8
	t,t-2,4-Nonadienal	20.8	20.8	20.8	20.8
	(2-Hydroxyethyl)-4-methyl-thiazol	947.8	947.8	947.8	947.8
	1-Octene-3-ol	0.8	0.6	0.6	0.6
15	1-Hexanol	3.6	3.5	3.6	3.6
	δ-Undecalactone	0.6	0.6	0.6	0.6
	2-Ethyl-1-hexanol	0.5	0.5	0.5	0.5
	2-Methyl-3-furanthiol	-	1.6	1.6	1.6
20	Methanedithiol	-	-	1.6	-
	Dithiomethane-diacetate	-	-	-	1.6
	Ethanol (up to 1000)	3.8	2.2	0.6	0.6

25 t=trans; c=cis

EPO - DG 1

BO 43104

17

30. 09. 1999

## CLAIMS

(59)

1. A flavoured foodstuff comprising an effective amount  
5 of at least one compound with a (hydrogenated) thio moiety  
and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a  
2-methyl-3-furyl-thio moiety and an effective amount of at  
least one compound having the structure

10  $U-CH_2-S-T$

in which C, H and S have the conventional meanings of  
carbon, hydrogen and sulphur atoms respectively, U  
represents a thiol group, a lower thioacyl group, a lower  
15 thioalkylgroup, a hydroxyl group or a 2-methyl-3-  
furyldithiogroup and T represents a hydrogen atom, a lower  
acyl group or a 2-methyl-3-furyl-thio group or a  $-S-CH_2-U$   
group as defined above.

20 2. A flavoured foodstuff according to claim 1, in which  
lower thioacyl- and lower acyl group means that these  
groups comprise from 2 to 6, preferably 2 or 3 carbon  
atoms.

25 3. A flavoured foodstuff according to claim 1 or 2, in  
which U represents a lower thioacyl group and T represents  
a lower acyl group.

4. A flavoured foodstuff according to a preceding claim  
30 in which lower thioacyl group means thioacetoxy and lower  
acyl group independently means acetyl.


5. A process for imparting a savoury flavour to a  
foodstuff comprising incorporating in said foodstuff an  
35 effective amount of at least one compound with a

BO 43104

18

(hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a 2-methyl-3-furyl moiety and an effective amount of at least one

5 compound having the structure



in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U

10 represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a lower acyl group or a 2-methyl-3-furyl-thio moiety or a  $-S-CH_2-U$  group as defined above.

15

6. A process according to claim 5, in which lower thioacyl-, lower alkyl- and lower acyl group means that these groups comprise from 2 to 6, preferably 2 or 3 carbon atoms.


20

7. A process according to claim 5 or 6 in which U represents a lower thioacyl group or a lower acyloxy group and T represents a lower acyl group.

25

8. A flavouring composition for foodstuffs comprising at least one compound comprising a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a 2-methyl-3-furyl-thio moiety and at least one compound having the structure

30



in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively,

35




BO 43104

19

U represents a thiol group, a lower thioacyl group, a hydroxyl group or a 2-methyl-3-furyldithio group and T represents a hydrogen atom, a lower alkyl group or a lower 5 acyl group.

9. A composition according to claim 8, in which lower thioacyl-, lower alkyl- and lower acyl group means that these groups comprise from 2 to 6, preferably 2 or 3 carbon 10 atoms.

10. The use of both an effective amount of at least one compound with a (hydrogenated) 2-methyl-3-furyl-thio moiety and a hydrogen atom, an  $-S-CH_3$  group, an  $-CO-CH_3$  group or a 15 (hydrogenated) 2-methyl-3-furyl-thio group and an effective amount of at least one compound having the structure



20 in which C, H and S have the conventional meanings of carbon, hydrogen and sulphur atoms respectively, U represents a thiol group, a lower thioacyl group, a lower thioalkylgroup, a hydroxyl group or a (hydrogenated) 2-methyl-3-furyldithio group and T represents a 25 hydrogen atom, a lower acyl group or a (hydrogenated) 2-methyl-3-furyl-thio group or a  $-S-CH_2-U$  group as defined above.

11. A process for preparing a pure compound with at 30 least one free thiol group as defined in claim 1 by hydrolysing the corresponding thioacyl compound in the presence of an enzyme or a cation exchange resin.

12. A process according to claim 12, in which the enzyme 35 is a lipase.



## Abstract

The invention relates to foodstuffs having a savoury  
5 flavour, to a process for their preparation and to  
compositions having a savoury flavour which products  
comprise (hydrogenated) 2-methyl-3-furanthiol and/or a  
derivative thereof and methanedithiol and/or a derivative  
thereof. Under derivatives are here to be understood their  
10 esters like thioacetates, their thioethers like methyl  
sulfides their disulphides etc. The combination of the  
above identified two groups of sulphur compounds in ppb  
levels leads to a foodstuffs with powerful flavour  
reminiscent of beef broth.

